

No 4  
47 South 8<sup>th</sup>

An  
Inaugural Essay On  
Injuries of the Head.

For

The Degree of Doctor of Medicine,  
In the University of Pennsylvania

By

Peyton R. Berkeley  
of Virginia.

Philadelphia

January 1828.



*[Faint, mirrored handwriting, likely bleed-through from the reverse side of the page. The text is illegible due to fading and mirroring.]*

by exte  
interna  
healthy  
The m  
must g  
of the  
T  
simila  
consta  
The da  
readu  
limab  
the p  
and th  
shoul



1  
By injuries of the head we mean such as are produced by external violence, consequently, not including those which arise from internal disease, or mal. conformation of the organs essential to the healthy functions of this important member of the human system. The most natural division of the subject is suggested by the arrangement of the parts most usually affected: viz: Injuries of the Scalp. of the Cranium; of the Membranes and of the brain itself—

Simple incised wounds of the scalp are to be treated as similar wounds in any other part; the surgeon, however, should constantly bear in mind the great necessity of immediate union. The danger of permitting suppuration in such cases can be readily perceived when we recollect the immediate and intimate connection existing between the periosteum and bone; the free interchange of vessels between the periosteum & Dura Mater; and the great probability that the brain must be affected, should these connections be by any means deranged. It is owing



to this  
so little  
injury  
ulation  
least  
this it  
and a  
ing to  
pretor  
so cony  
nosis, a  
perfect  
of thos.  
Pun  
true to  
striking  
be attr  
are ma  
the bon



to this connection that even an experienced surgeon can with so little certainty foretell the termination of the most superficial injury of the head. To day all may appear well, healthy granulations, wound maturing well, patient free from pain; let the least transfer of irritation to the D.M. be produced (and to effect this it is only necessary that the connecting vessels be disturbed) and all these favourable symptoms vanish, and that without warning to the surgeon by which such a change might have been foretold. Certain it is, that there is no branch in surgery which so completely baffles the skill of the surgeon to give a correct prognosis, the most horrid injuries frequently resulting in the most perfect cure; and the reverse being often the sad termination of those which at first scarcely excited attention.

Punctured wounds are much the most dangerous. This remark, true to a certain extent, as respects the whole system, is most strikingly exemplified in wounds of the scalp. This we think may be attributed to two causes 1<sup>st</sup> The weapon, with which such wounds are made, from its construction, is usually driven directly against the bone, and consequently is more liable to injure the cranium



than  
off the  
it be  
thus  
of the  
dis-cry  
of injur  
win a  
even  
throug  
the m  
in ma  
not ex  
appear  
injur  
"the sk  
small  
pale fo  
not ver



than when it is of such a nature as only to cut through or pare off the scalp. 2<sup>d</sup> The wound is small and deep, the matter, should it be formed, has no sufficient aperture for its escape, and being thus confined is more liable to affect the interchanged vessels of the Periosteum and D. M. than in wounds of almost any other description. Hence the size of a wound is no evidence of the extent of injury. Various other considerations are to be taken in to form even a probable conclusion of what may be the result. Punctured wounds do not always extend entirely to the bone, but only through particular portions of the scalp, as *cutis Vera*, tendons of the muscles &c. According to Pott, the effects of the injury will in many cases indicate the extent of injury. Thus should the wound not extend to the Aponeurosis: Expansion, the unfavourable appearances, should any follow, will not be confined to the part injured, but will include the head and face. In his own language, "the skin wears a yellowish tint, and is sometimes thick set with small blisters; it receives the impression of the fingers, and becomes pale for a moment, but soon returns to its inflamed colour; it is not very painful to the touch, and the eyelids and ears are al-



... of such a nature as may be out of the way of the  
... the means is to make the matter more  
... it is not sufficient to make the matter more  
... is more likely to effect the intended change  
... and it is then in a measure of a more  
... the use of a variety of means of the same  
... these considerations are to be taken as the basis  
... of which may be the most beneficial  
... not only to the health of the body but also  
... of the body as well as the mind  
... to the effect of the remedy  
... the extent of the remedy  
... the most common  
... and not be confined to the first  
... the body and face to his own  
... and is a matter which is not  
... of the finger and thumb  
... of the body and face  
... and the effect of the remedy

ways  
time  
short  
pulse  
ding  
found  
ger  
is ent  
thoug  
which  
grow  
ant  
after  
is slo  
violet  
the h  
cold  
and  
The



4.  
ways included in the tumefaction, the former of which are sometimes so much distended as to be closed; a febrile heat and thirst generally accompany it. the patient is restless. has a quick pulse. most commonly nausea and inclination to vomit. According to that able writer, this affection, as just described, is only found in bilious habits; by one not accustomed to it, great danger might be apprehended; but Pott assures us this apprehension is entirely unfounded. It is an inflammatory affection, and although partaking in some degree of the Erysipelatous Affection which frequently follows injury of the head, yet it is less dangerous, and may be distinguished. In the first place, that dependent on a bilious habit, makes its appearance in a short time after the injury is inflicted, one or two days: whereas Erysipelas is slow and gradual; the former is comparatively mild; the latter violent; a quick, hard, ~~stord~~ pulse; excessive heat; violent pain in the head; extreme anxiety and restlessness; and not unfrequently cold shiverings. In both species the wound has an unhealthy appearance, but in the latter the scalp is separated from the cranium. The treatment of this more simple Affection is very simple.



Rest,  
pholo  
all the  
Where  
proves  
is often  
prevent  
practi  
has been  
tumors  
commen  
compos  
yet wh  
The in  
placed  
two we  
the scap  
prefer  
to the



Purgatives, Blisters, venesection, in other words the antiphlogistic treatment, regulated according to circumstances, is all that is required. This simple treatment will often effect a cure.

Where the system is in a phlogosed state, and the inflammation<sup>m</sup> proves obstinate, the division of the scalp is recommended, and is often a means of expediting the cure. This will generally prevent sloughing, and consequently diminish danger. This practice, Division of the scalp, is also recommended where blood has been extravasated under the scalp, forming a small round tumour. The most simple treatment of such Tumours is that recommended by Mr. S. Cooper. It consists in mild purges, and a lotion composed of *Acidum Aceticum* (Vinegar) and *Muriate of Ammonia*. Yet when this fails, the incision must be resorted to.

The indication in all incised wounds is the same. The parts are to be placed in exact apposition, and thus retained. This can be effected in two ways. by the interrupted suture or adhesive straps. For wounds of the scalp the straps are vastly preferable, and should always be preferred, where they will answer the purpose intended. To apply them to the head, it is necessary that it be previously shaved. The straps



a satisfaction  
simple  
With  
portions  
greatly  
genera  
every e  
rated, a  
other co  
ity, an  
of the p  
fear u  
formed  
evanc  
The int  
ing left  
It has  
stances,  
satisfac



a suture, with the antiphlogistic regimen, is as much as simple incised wounds of the scalp generally demand.

With regard to the treatment of those cases where considerable portions of the scalp have been torn up, surgeons of eminence once greatly differed, although, at present, we believe, no fact is more generally admitted than that the part should be replaced, and every exertion made to preserve it. Unless the flap be entirely separated, a surgeon can never be justifiable in pursuing any other course. It is undoubtedly correct, since it prevents deformity, and in no respect militates against the speedy recovery of the patient. In many cases where such an attempt would appear useless, union is accomplished, even where matter has been formed beneath the flap requiring several perforations for its evacuation.

The entire removal of the scalp, the bone immediately beneath being left bare, is sometimes the effect of a severe blow on the cranium. It has often occurred to me to examine whether under such circumstances, exfoliation be a necessary consequence. This question has been satisfactorily answered by a short note in Pott. He says it depends



on other  
Persians  
repels, a  
cipally a  
ular, a  
of the bon  
ing off  
in the co  
either by  
of art, is  
that fra  
be carri  
which m  
Fruct  
many spe  
semblan  
surgeons  
clapped  
depression



on other circumstances besides the mere removal of the scalp and Periosteum. The solidity of the surface of the bone, the size of the vessels, and the impulse of the blood through them, are what principally determine it. If the bone be favourable in these particulars, a granulation of flesh will be generated on the surface of the bone, which will cover and firmly adhere to it, without throwing off the smallest exfoliation, especially in young subjects. On the contrary if the circulation is interrupted in the bone, either by its natural density and hardness, or by the applications of art, it must part with a scale to a certain depth, that is that part of the surface, through which the circulation ceases to be carried on, will be separated from and cut off by the vessels which nourish the rest of the bone.

Fractures of the Cranium were by the old writers divided into many species, the name of each being derived from a supposed resemblance of the fissure or fracture to some familiar object. Modern surgeons have rejected this long and difficult list, and instead classed the whole under two heads viz: fractures with & without depression; and these include all possible cases. It was a practice



the no more than to show the same as if the  
the activity of the surface of the bone the size of the  
and the shape of the bone through them, and what you  
may observe it is the bone to be examined in this place.  
In a preparation of this sort the bones are in the shape  
of the bone in all cases and finally when it is finished the  
of the smallest preparation is usually in a very simple  
the surface of the preparation is interrupted in the bone.  
of the natural shape and shape, as if the preparation  
it must have with a view to a certain effect. That is  
the part of the surface, through which the preparation is made  
may be made to separate from and cut off by the surface  
of the bone, the part of the bone.  
The preparation of the bone is made by the use of the bone  
and the bone of each being covered from a surface of the  
of the bone a preparation to some extent of the bone.  
The bone is prepared in this way and without any  
the bone is prepared in this way and without any  
the bone is prepared in this way and without any

with the  
skull;  
figure  
the  
with  
this is  
object  
accomplish  
get it  
is new  
the  
tion of  
and  
form  
figure  
the  
ing the  
really  
mischa



with the old operators to use the Trepan in every case of fractured skull; several perforations were usually made along a simple fissure, and this before a symptom of compression was manifest, with the ostensible purpose of preventing the bad effects which sometimes result from such injuries. Fortunately this useless and painful practice is now abandoned. Experience has undoubtedly proved that the object of these repeated perforations is as certainly, and more certainly, accomplished by a rigorous adherence to the Antiphlogistic Treatment. Yet it is not to be understood from these remarks that the trepan is never demanded in cases of simple fissure. Even Abernethy (and the opinion of no one more merits our attention) admits, "the operation of the trepan is frequently performed in cases of simple fracture, and that very judiciously and properly; but it is not performed because the bone is broken or cracked; a crack or fissure of the skull can never require perforations; or that the Dura Mater under it be laid bare; the reason for doing this springs from other causes than the fracture, and these really independent of it; they spring from the nature of the mischief which the parts within the cranium have sustained, and



not  
all the  
inde  
rule of  
author  
than  
ularly  
case of  
result,  
without  
recove  
Many  
duce  
be be  
fortune  
myself  
the ope  
pursue  
requir



not from the accidental division of the bone. From these arise all the threatening symptoms, and from these the necessity & vindication of performing the operation of the trepan." This rule for applying the Trepan, according to the same high authority, should guide in cases of depressed fracture, not less than in those of fracture without depression. Bad effects, particularly compression of the brain, are to be apprehended in every case of depressed fracture. Yet such is not uniformly the result, in as much as many instances are on record, where, without assistance from the surgeon, patients thus injured have recovered, and without permanent injury or deformity. Many cases, illustrative of the position assumed, might be adduced, but the fact is now so generally admitted that it would be but waste of time, both to the professor who may be so unfortunate as to be compelled by duty to peruse this piece, and to myself. In every instance then the surgeon, in determining on the operation, must be directed by circumstances; we are persuaded that very few cases, comparatively speaking, will require the instrument. I have had the good fortune during the



last s  
in w  
me, in  
thing,  
fell d  
few ca  
uses th  
in the  
were  
the op  
(We sh  
of op  
From  
are p  
often  
the m  
jury  
succ  
and




last summer, of attending several cases of fractured skull, and in every instance a speedy recovery was the result, except in one, in which case my preceptor performed the operation. Nothing, however, could have given relief in this case, as the child fell directly on its head, from the third story of a high house. These few cases, and the uniform success of my preceptor, who seldom uses the trepan, may have biased my mind; yet I feel confident in the correctness of the position assumed, that the old operators were too hasty in the employment of this instrument, and that the operation is not demanded in every case of depressed fracture. We shall not, as we deem it unnecessary, describe the manner of operating with the Trepan.

From a slight blow or fall, even where no marks of violence are perceptible on the scalp, the most serious consequences have often resulted. The first indication of a transfer of irritation to the membranes or brain, is pain under the part where the injury has been received. This gradually increases; high fever succeeds; pulse is quick and hard; skin hot; patient restless; and if no remedies be used to allay the inflammation, the



*[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]*



*[Faint, illegible handwriting on the right edge of the page, likely bleed-through from the reverse side.]*



symptoms increase and in a short time become violent; the patient is kept constantly awake, or if asleep, it is broken and unrefreshing; the pain extends over the whole head. Should a wounded scalp be the precursor, about this time the wound changes its appearance; discharges a thin serum, instead of healthy pus, and the lint adheres. Should the Periosteum be in sight, it will be found to have changed its colour; it will be darker than usual; and be found separated from the bone. Such symptoms denote the separation of the Dura Mater from the Cranium, and the formation of matter between, and generally result in evident symptoms of compression. The injury is sometimes even more extensive. The inflammation may extend to the other membranes, and the brain, and then instead of matter under the Cranium upon the D. M. it may be beneath it; or between the P. M. and brain; or in the ventricles of the brain. All this train of evils is the effect of inflammation, produced by external violence, and yet without any outward mark. In other cases the violence may be so great as to rupture the blood vessels immediately. Then there will be an effusion of blood, and compression the probable result. Compression



wh  
or  
sign  
puls  
light  
and  
plic  
app  
geon  
wre  
the  
seve  
lefs  
fora  
ed.  
ion,  
pur  
these  
The



whether produced by depression of bone, the formation of matter, or the ~~eff~~ extravasation of blood, is characterised by the following symptoms "The pulse will be found slow and regular; the pupils of the eyes greatly dilated, and insensible to the strongest light; the breathing stertorous, slow and difficult; the limbs loose, and yielding, and sometimes paralytic; the insensibility complete, so much so that the patient cannot be roused by the application of the strongest stimulants." Had one of the old Surgeons been called to a patient in this situation, not a moment would have been lost. The trepan, the trepan would have been the reiterated call; nor would they have hesitated to make several perforations in the course of the same fracture. Doubtless this procedure in a majority of cases (I mean a single perforation) would be correct, but frequently entirely unwarranted. According to some of the highest authorities in our profession, the Trepan should not be first employed. Venesection, mild purges (when the case will warrant delay) are the first steps. If these fail to restore, the propriety of the operation is unquestionable. The point now to be determined is, where to apply the instrument.



If a  
ed.  
ma  
ton  
sore  
stan  
If p  
ed.  
the  
tens  
thes  
will  
ever  
some  
for t  
is sc  
prep  
light  
bled



If a wound or bruise or any external mark exists, it is easily determined. But where no such land mark is discoverable, it becomes a matter of the greatest doubt and uncertainty. Should the symptoms be urgent, no time is to be lost, the operation must be resorted to. There are considerations, which, under such circumstances, should have their influence in directing the Surgeon. If pain is felt from pressure on any part, a useful hint is suggested. The kind of instrument by which the injury was inflicted; the relative position of the two individuals &c. &c. Or if the patient has fallen, the manner of falling; the direction &c. &c. all these should be carefully examined, and we are persuaded they will seldom fail to give some useful hint. Should they, however, and the operation be positively demanded, let it be over some large blood vessel, as the middle artery of the D. M. and for this reason, the probability of extravasation here is greatest. It is scarcely necessary to remark, that after the perforation, the depressed bone should be elevated, blood or matter evacuated; wound lightly dressed; patient confined to the antiphlogistic regimen, and bled and purged *pro re nata*. for four or five weeks. &c. &c.



The first of these is the fact that the  
the second is the fact that the  
the third is the fact that the  
the fourth is the fact that the  
the fifth is the fact that the  
the sixth is the fact that the  
the seventh is the fact that the  
the eighth is the fact that the  
the ninth is the fact that the  
the tenth is the fact that the  
the eleventh is the fact that the  
the twelfth is the fact that the  
the thirteenth is the fact that the  
the fourteenth is the fact that the  
the fifteenth is the fact that the  
the sixteenth is the fact that the  
the seventeenth is the fact that the  
the eighteenth is the fact that the  
the nineteenth is the fact that the  
the twentieth is the fact that the  
the twenty-first is the fact that the  
the twenty-second is the fact that the  
the twenty-third is the fact that the  
the twenty-fourth is the fact that the  
the twenty-fifth is the fact that the  
the twenty-sixth is the fact that the  
the twenty-seventh is the fact that the  
the twenty-eighth is the fact that the  
the twenty-ninth is the fact that the  
the thirtieth is the fact that the  
the thirty-first is the fact that the  
the thirty-second is the fact that the  
the thirty-third is the fact that the  
the thirty-fourth is the fact that the  
the thirty-fifth is the fact that the  
the thirty-sixth is the fact that the  
the thirty-seventh is the fact that the  
the thirty-eighth is the fact that the  
the thirty-ninth is the fact that the  
the fortieth is the fact that the  
the forty-first is the fact that the  
the forty-second is the fact that the  
the forty-third is the fact that the  
the forty-fourth is the fact that the  
the forty-fifth is the fact that the  
the forty-sixth is the fact that the  
the forty-seventh is the fact that the  
the forty-eighth is the fact that the  
the forty-ninth is the fact that the  
the fiftieth is the fact that the  
the fifty-first is the fact that the  
the fifty-second is the fact that the  
the fifty-third is the fact that the  
the fifty-fourth is the fact that the  
the fifty-fifth is the fact that the  
the fifty-sixth is the fact that the  
the fifty-seventh is the fact that the  
the fifty-eighth is the fact that the  
the fifty-ninth is the fact that the  
the sixtieth is the fact that the  
the sixty-first is the fact that the  
the sixty-second is the fact that the  
the sixty-third is the fact that the  
the sixty-fourth is the fact that the  
the sixty-fifth is the fact that the  
the sixty-sixth is the fact that the  
the sixty-seventh is the fact that the  
the sixty-eighth is the fact that the  
the sixty-ninth is the fact that the  
the seventieth is the fact that the  
the seventy-first is the fact that the  
the seventy-second is the fact that the  
the seventy-third is the fact that the  
the seventy-fourth is the fact that the  
the seventy-fifth is the fact that the  
the seventy-sixth is the fact that the  
the seventy-seventh is the fact that the  
the seventy-eighth is the fact that the  
the seventy-ninth is the fact that the  
the eightieth is the fact that the  
the eighty-first is the fact that the  
the eighty-second is the fact that the  
the eighty-third is the fact that the  
the eighty-fourth is the fact that the  
the eighty-fifth is the fact that the  
the eighty-sixth is the fact that the  
the eighty-seventh is the fact that the  
the eighty-eighth is the fact that the  
the eighty-ninth is the fact that the  
the ninetieth is the fact that the  
the ninety-first is the fact that the  
the ninety-second is the fact that the  
the ninety-third is the fact that the  
the ninety-fourth is the fact that the  
the ninety-fifth is the fact that the  
the ninety-sixth is the fact that the  
the ninety-seventh is the fact that the  
the ninety-eighth is the fact that the  
the ninety-ninth is the fact that the  
the hundredth is the fact that the

long  
ing  
love  
occa  
tree  
beg  
ing  
from  
inj  
or  
sho  
cou  
wou  
com  
at  
he  
lou  
lat  
bth



Sometimes after a perforation, the brain protrudes through the opening, and gradually increasing produces what is styled *Fungus cerebri*. The treatment for such a tumour is light dressings, with occasionally light moderate pressure.

Even after the removal of a portion of the bone, the surgeon may be greatly embarrassed. The D. M. may be perfectly clean, exhibiting no sign of blood or effused pus; it may not even be detached from the cranium. Thus situated he should remember that the injury may be deeper seated; the fluid may be under the D. M. or still farther from the surface. Here the question would arise should this membrane be perforated. Where no hopes of recovery could be indulged unless the brain be immediately relieved, no one would hesitate to proceed. But the case should be a desperate one. The common method is merely to puncture this membrane with a lancet or other sharp instrument. Sir A. Cooper, however, remarks, that he has never known a single case of recovery where the D. M. alone has been punctured. But strange as it may appear, he has related several cases of complete recovery where the D. M. & P. M. were both divided. Accordingly he follows this practice, and strenuously



address  
a su  
may  
sist  
to her  
two.  
must  
throu,  
shou  
liable  
and  
this  
the ly  
Comp  
person  
lang  
These  
time  
under



advises it. In attending a lecture of Dr. Hopkinson, very lately, I heard a suggestion from him, which, so far as I know, is original. be this as it may, it appeared to me to merit attention. The D. M. is known to consist of two laminae or layers, an internal and an external. It occurred to him, that the puncture ought not to be made directly through the two, but that the external coat should be first divided, the instrument then so directed as to pass between the two a short distance, and then through the internal. The orifice could be easily dilated until any fluid should be discharged. He thinks that inflammation will not be more liable to follow such a puncture, than the one commonly resorted to, and fungus cerebri not so likely to occur. Yet, however performed, this is a dangerous operation, and should not be resorted to, unless the life of the patient is in immediate danger.

Compression is not always the result of severe blows on the head. The person is more frequently only stunned, as it is stated in familiar language; or in medical language he suffers a concussion. These two affections, Compression & Concussion are sometimes so intimately blended in the same case, that it is difficult to determine under which the patient labours. Thus a person will receive a severe



the following a list of the symptoms of the disease  
as far as I have been able to ascertain. It is  
a disease of the lungs, and is attended with  
a cough, which is at first dry, but soon  
becomes loaded with phlegm. The expectoration  
is at first thin and watery, but soon  
becomes thick and yellow. The patient  
is attended with a shortness of breath,  
and a pain in the chest, which is  
increased by the cough. The pulse is  
at first small and frequent, but soon  
becomes full and slow. The disease  
is attended with a loss of appetite,  
and a general debility. The patient  
is attended with a fever, which is  
at first low, but soon becomes high.  
The disease is attended with a  
cough, which is at first dry, but soon  
becomes loaded with phlegm. The expectoration  
is at first thin and watery, but soon  
becomes thick and yellow. The patient  
is attended with a shortness of breath,  
and a pain in the chest, which is  
increased by the cough. The pulse is  
at first small and frequent, but soon  
becomes full and slow. The disease  
is attended with a loss of appetite,  
and a general debility. The patient  
is attended with a fever, which is  
at first low, but soon becomes high.

(low)  
nupt  
con  
on co  
a cas  
are,  
tus; p  
blow  
first  
extre  
ing; n  
ishd.  
and  
lated  
puls  
the p  
with  
the la  
and



blow, symptoms of concussion appear; by the violence a vessel is ruptured; by the time the system has recovered from the effects of concussion, a sufficient quantity of blood is extravasated to bring on compression, the patient being all the while insensible. In such a case great doubt must exist. The symptoms of simple concussion are, Trembling; vertigo; sickness of stomach; loss of mental faculties; partial dimness of sight. According to Mr S. Cooper when the blow has been very severe, the following will be the symptoms. At first the patient is stunned and in a state of total insensibility; the extremities are frequently cold, his pulse weak, slow, and intermitting; his respiration hardly perceptible; his powers of motion abolished. The following symptoms then will distinguish concussion and compression. In the former the pupil is either moderately dilated or contracted; in the latter greatly dilated; in the first the pulse is weak and tremulous; in the latter slow and regular; the former without stertorous breathing; the latter always attended with it; the former is always accompanied with sickness of stomach; the latter never. In some cases these will be variously combined, and the true nature of the injury difficult to determine.



The  
The  
to the  
erally  
soon  
infl  
We  
cise  
rest.  
collec  
ject.  
purly  
style, a  
in the  
case  
duct  
loose a  
from  
must



The treatment in cases of concussion next claims our attention. The antiphlogistic regimen, mild purgatives; cold applications to the head, after shaving it; keeping the head elevated, will generally be sufficient. We should be very cautious not to bleed too soon; not until reaction is established. we then bleed to keep down inflammation.

We have endeavoured to give our ideas on this subject in a concise manner. I am conscious that this paper can afford but little interest. The sentiments are stale and hackneyed. Yet it should be recollected that without experience, no man can write on such a subject. We have intentionally omitted much, which might have been properly introduced. I have aimed particularly at brevity, sacrificing ease of style, and sometimes more fear, perspicuity, to attain it. We will conclude in the words of Mr. Pott, "the peculiar circumstances of each individual case must furnish direction to the surgeon for his particular conduct. The parts which are depressed must be elevated; such as are loose and cannot be brought to lie even, such as cannot be prevented from pressing on the membranes, or such as wound and irritate it, must at all events be taken away. The free discharge of blood and



ly my  
these  
earn  
apis



lymph at present, and matter in future, must be provided for, and therefore every symptom and appearance must be carefully and earnestly attended to, least the most proper <sup>or</sup> opportunity of giving assistance be not embraced. *Itatiam! Itatiam!* [F.L.]

*Receipts employed by me*

*in supplying the*

*for the*

*of Doctor of Medicine*

*in the*

*University of Pennsylvania*

*By*

*James M. Smith*

*South Carolina*



10.  
The first and most important  
principle of the system is  
that the mind must be  
trained to receive and  
retain impressions.